

## The Center of Evidence References

Badke M, Shea T, Miedaner J, et al. Outcomes After Rehabilitation for Adults with Balance Dysfunction. *APMR*. 2004; 85(2):227-233.

Dite W, Temple VA. A clinical test of stepping and change of direction to identify multiple falling older adults. *Arch Phys Med Rehabil*. 2002;83:1566-71.

Fansler C, Poff C, Shepard K, et al. Effects of Mental Practice on Balance in Elderly Women. *PT*. 1985; 65(9):1332-1338.

Hanke T, Tiberio D. Lateral Rhythmic Unipedal Stepping in Younger Middle Aged and Older adults. *J of Ger PT*. 2006;29(1): 20-25.

Kobayashi R, Hosoda M, Minematsu A, et al. Effects of Toe Grasp Training for the Aged on Spontaneous Postural Sway; *J PT SCI*. 1999; 31(1):31-34.

Kobayashi R, Hosoda M, Minematsu A, et al. Effects of Toe Grasp Training for the Aged on Spontaneous Postural Sway; *J PT SCI*. 1999;31(1):31-34

Li F, Harmer P, Fisher J, et al. Tai chi and Fall Reduction in Older Adults: A Randomized Controlled Trial. *J of Gero*. 2005;60A(2): 187-194.

Sakamoto J, Nakamura T, Hagino H, et al. Effects of unipedal standing balance exercise on the prevention of falls and hip fracture among clinically defined high-risk elderly individuals: a randomized controlled trial. *Orthop Sci*. 2006;11:467-472.

Sakamoto J, Nakamura T, Hagino H, et al. Effects of unipedal standing balance exercise on the prevention of falls and hip fracture among clinically defined high-risk elderly individuals: a randomized controlled trial. *Orthop Sci*. 2006;11:467-472

Waddington G, Adams R. The Effect of a 5-Week Wobble-Board Exercise Intervention on Ability to Discriminate Different Degrees of Ankle Motion Inversion, Barefoot and Wearing Shoes: A Study in Healthy Elderly. *JAGS*. 2004; 52(4): 573-576.

Yardley L, Donovan-Hall M, Smith H, et al. Effectiveness of Primary Care Based Vestibular Rehabilitation for Chronic Dizziness; *Ann Int Med*. 2004; 141(8):598-605.

Feland J, Myrer J, Schulthies S, et al. The Effect of Duration of Stretching of the Hamstring Muscle Group for Increasing Range of Motion in People Aged 65 Years or Older. *PT*. 2001; 81(5):1100-1117.

Hauer K, Rost B, Rutschle K, et al. Exercise Training Rehabilitation and Secondary Prevention of Falls in Geriatric Patients with a History of Injurious Falls. *JAGS*. 2001; 49(1):10-20.

Kerrigan D, Xenopoulous-Oddsson A, Sullivan M, et al. Effects of a Hip Flexor-Stretching Program on Gait in the Elderly. *APMR*. 2003; 84(1):1-6.

Mecagni C, Smith J, Roberts K, et al. Balance and Ankle Range of Motion in Community-Dwelling Women Aged 64 to 87: A Correlational Study. *PT*. 2000; 80 (10):1004-1011.

Moreside J, Vera-Garcia F, McGill S. Trunk Muscle Activation Patterns, Lumbar Compressive Forces, and Spine Stability When Using the Bodyblade. *PT*. 2007; 87(20):153-163.

Schiller, J, Schubert, S, Lowe, S ;The Effects of a 12-week Progressive Resistance Program on Lower Extremity Strength and Timed Up & Go Measures in Community Dwelling Elders: A Pilot Study. *Gerinotes*. 2001; 8(6):12-20.

Campbell J, Robertson M, Gardner M, et al. Randomized controlled trial of general practice program for home based exercise to prevent falls in elderly women. *BMJ*. Oct 25,1987; Vol 315: 1065-69.

Ledin T, Kronhed A, Moller C, et al. Effects of Balance Training in Elderly Evaluated By Clinical Tests and Dynamic Posturography. *J Vest Res*. 1991;1(2):129-138.

Nelson M. The effects of multidimensional Home-based exercise on functional performance in elderly people. *Journal of Gerontology*. 2004; 59: 154-160.

Nishimoto J, Nakamura S, Imai T, et al. Increasing Gait Performance in Elderly Women Using a Step Exercise Program. *J of PT Sci*. 1999;11(2):71-78.

Nitz, J, Choy, N. The Efficacy of a specific balance-strategy training program for preventing falls among older people. *Age & Aging*. 2004; 33(1):52-58.

Reece A, Simpson J. Preparing Older People to Cope after a Fall. *Physiotherapy*. April 1996; 82 (4): 227-235.

Sakamoto J, Nakamura T, Hagino H, et al. Effects of unipedal standing balance exercise on the prevention of falls and hip fracture among clinically defined high-risk elderly individuals: a randomized controlled trial. *Orthop Sci*. 2006;11:467-472.

Toulotte C, Fabre C, Dangremont B, et al. Effects of physical training on the physical capacity of frail, demented patients with a history of falling: a RCT. *Age and Ageing*. 2003; 32(1):67-73.

Fitzgerald G, Childs J, Ridge T, et al. Agility and Perturbation Training for Physically Active Individual With Knee Osteoarthritis. *PT*. 2002; 82(40): 372-382.

Gardner M, Robertson M, McGee R, et al. Application of a Falls Prevention Program for Older People to Primary Health Care Practice. *Prev Med.* 2002; 34: 546-553.

Ledin T, Kronhed A, Moller C, et al. Effects of Balance Training in Elderly Evaluated By Clinical Tests and Dynamic Posturography. *J Vest Res.* 1991; 1(2): 129-138.

Li F, Harmer P, Wilson N, et al. Health Benefits of Cobblestone Mat Walking; *J of Aging and Phys Act.* 2003; 11: 487-501.

Robertson M, Campbell A, Gardner M, et al. Preventing Injuries in Older People by Preventing Falls: a Meta- Analysis of Individual Level Data. *JAGS.* 2002; 50(5):905-911.

Sidaway G, Anderson J, Danielson G, et al. Effects of Long-Term Gait Training Using Visual Cues in Individuals with Parkinson's Disease. *PT.* 2006; 86(2): 186-194.

Silsupadol P, Siu K, Shumway-Cook A, et al. Training of Balance Under Single and Dual Task Conditions in Older Adults With Balance Impairments. *PT.* 2006; 86(2): 269-281.